

Maximum residue level for cyantraniliprole


Published by AGRINFO on 29 Jun 2023

For more information see the [full record](#) on the AGRINFO website – where you can also view the latest [AGRINFO Update](#) newsletters and [search](#) the database.

Tables & Figures

Table 1 Changes to maximum residue levels for cyantraniliprole			
Food category	Products	Cyantraniliprole (mg/kg)	
		Old MRL	New MRL
Stone fruits	Apricots	0.01*	0.7
Root and tuber vegetables	Potatoes, cassava roots/ manioc, sweet potatoes, yams, arrowroots	0.05	0.15
Fruiting vegetables	Melons, pumpkins, watermelons	0.3	0.4
Brassica vegetables	Chinese cabbages/ pe-tsai	0.01	30
Lettuce and salad plants	Lamb's lettuces/ corn salads, escaroles/ broad-leaved endives, cresses and other sprouts and shoots, land cresses, Roman rocket/ rucola, red mustards, baby leaf crops	0.01	15
	Lettuces	5	15
Spinaches and similar leaves	Purslanes, chards/ beet leaves	0.01*	20
Herbs and edible flowers	Parsley	0.02*	20
Oilseeds	Linseeds; seeds of poppy, sesame, mustard, pumpkin, safflower, borage, gold of pleasure, hemp; castor beans	0.01*	1.5

* Limit of determination.



 www.agrinfo.eu

Source: based on Regulation [2023/1068](#)

Disclaimer: *Under no circumstances shall COLEAD be liable for any loss, damage, liability or expense incurred or suffered that is claimed to have resulted from the use of information available on this website or any link to external sites. The use of the website is at the user's sole risk and responsibility. This information platform was created and maintained with the financial support of the European Union. Its contents do not, however, reflect the views of the European Union.*