

Maximum residue level for propamocarb

Published by AGRINFO on 04 Jun 2024; Revised 12 Dec 2024

[Draft](#) Commission Regulation as regards maximum residue levels for dimoxystrobin, ethephon and propamocarb in or on certain products

[Draft](#) Annex (PLAN/2024/1305 R2 DRAFT)

Commission Regulation (EU) [2024/1439](#) as regards maximum residue levels for fenazaquin, mepiquat and propamocarb in or on certain products

What is changing and why?

The EU proposes to lower the maximum residue level (MRL) for propamocarb on lettuces from 40 to 20 mg/kg, following concerns about the safety of the current MRL.

Earlier in 2024, the EU raised the MRL for propamocarb on honey and other apiculture products from 0.05 to 15 mg/kg.

Actions

Feedback on the EU's proposal ([G/SPS/N/EU/801](#)) closed on 3 February 2025.

Suppliers of lettuces should review their use of propamocarb and assess whether any changes will be needed to existing good agricultural practices (GAP) to ensure compliance with the new MRL.

Timeline

The new MRL on lettuces is expected to apply from approximately **January 2025**.

For more information see the [full record](#) on the AGRINFO website – where you can also view the latest [AGRINFO Update](#) newsletters and [search](#) the database.

Disclaimer: *Under no circumstances shall COLEAD be liable for any loss, damage, liability or expense incurred or suffered that is claimed to have resulted from the use of information available on this website or any link to external sites. The use of the website is at the user's sole*

risk and responsibility. This information platform was created and maintained with the financial support of the European Union. Its contents do not, however, reflect the views of the European Union.