

Maximum residue levels for acequinocyl

Published by AGRINFO on 04 Feb 2026; Revised 24 Jun 2026

[Draft](#) Commission Regulation as regards maximum residue levels for acequinocyl, deltamethrin, dodine, maleic hydrazide, pinoxaden and prothioconazole in or on certain products

[Draft](#) Annex [PLAN/2025/3053]

Commission Regulation (EU) [2026/140](#) as regards maximum residue levels for acequinocyl, chlormequat, metalaxyl-M, pyraclostrobin, sulfoxaflor and trifloxystrobin in or on certain products

What is changing and why?

The European Union (EU) is discussing increasing the maximum residue levels (MRLs) for acequinocyl on blueberries, cranberries, currants, and gooseberries from the limit of determination (LOD) of 0.01 mg/kg to 2 mg/kg. (The LOD is the lowest level that can be detected using the most modern and reliable analytical methods.) An application for a change of MRLs by the producer of acequinocyl was evaluated by the European Food Safety Authority, and the requested changes were found to be safe for consumers.

In February 2026, the EU increased the MRL for acequinocyl on strawberries from 0.01 (LOD) to 0.3 mg/kg.

Timeline

The new MRLs on blueberries, cranberries, currants, and gooseberries are expected to apply in the second half of 2026.

The new MRL on strawberries applies from **11 February 2026**.

For more information see the [full record](#) on the AGRINFO website – where you can also view the latest [AGRINFO Update](#) newsletters and [search](#) the database.

Disclaimer: *Under no circumstances shall COLEAD be liable for any loss, damage, liability or expense incurred or suffered that is claimed to have resulted from the use of information available on this website or any link to external sites. The use of the website is at the user's sole risk and responsibility. This information platform was created and maintained with the financial support of the European Union. Its contents do not, however, reflect the views of the European*

