

Maximum residue levels for cycloxydim



Published by AGRINFO on 30 Nov 2022

For more information see the [full record](#) on the AGRINFO website – where you can also view the latest [AGRINFO Update](#) newsletters and [search](#) the database.

Tables & Figures

Table 1 Maximum residue levels for cycloxydim				
Food category	Products	Cycloxydim (mg/kg)		
		Old MRL	New MRL	
Pome fruits	Apples, pears	0.1	0.09*	
Stone fruits	Apricots, peaches	0.2	0.09*	
Berries and small fruits	Table/wine grapes	0.5	0.4	
Miscellaneous fruits	Mangoes	0.2	0.09*	
Root and tuber vegetables	Potatoes	3	4	
	Cassava root/manioc/yams/arrowroots	0.2	0.09*	
	Sweet potatoes	0.2	0.6	
	Beetroots, horseradishes, parsnips	0.9	3	
	Celeriacs/turnip rooted	5	3	
	Parsley roots/Hamburg roots, Swedes/rutabagas	0.2	1	
	Radishes	0.2	3	
	Salsifies	1.5	3	
	Turnips	1	3	
	Bulb vegetables	Garlic, shallots	1	1.5
		Spring onion/green onions and other bulb vegetables	1	0.3
Brassica vegetables	Broccoli, kohlrabi	2	9	
	Cauliflowers, head cabbages	5	9	
	Brussel sprouts	6	9	
Leaf vegetables, herbs and edible flowers	Lamb's lettuces/corn salads	0.5	2	
	Cresses and other sprouts and shoots, land cresses	0.5	1	
	Roman rocket/rucola	0.5	3	
	Red mustards, baby leaf crops	0.2	3	
	Purslanes, chards/beet leaves	0.6	2	
	Grape leaves and similar, watercresses, witloofs/Belgium endives	0.2	0.09*	
	Chervil, parsley, sage, rosemary, thyme, basil and edible flowers, laurel/bay leaves, tarragon	0.2	2	
	Chives	0.2	0.8	
	Celery leaves	1	2	
	Legume vegetables	Beans (without pods)	2	15
Peas (with pods)		5	2	
Lentils		1	15	
Stem vegetables	Asparagus, cardoons, Florence fennels, rhubarbs, bamboo shoots, palm hearts	0.2	0.09*	
	Globe artichokes	2	0.09*	
	Celeriacs	1	0.09*	
Pulses	Lentils, lupins/lupini beans	5	20	
Oilseeds	Peanuts/groundnuts, seeds (sesame, pumpkin, safflower, gold of pleasure, hemp), castor beans	0.2	0.09*	
	Poppy seeds, borage seeds	0.2	7	
	Sunflower seeds	6	7	
	Rapeseeds/canola seeds	9	7	
	Mustard seeds	5	6	
	Cotton seeds	0.5	7	
	Herbal infusions from	Valerian, ginseng	7	9
Sugar plants	Sugar beet roots	0.5	0.2	
Animal products: bovine & equine & other farmed terrestrial animals	Muscle	0.06	0.1	
	Fat	0.01	0.15	
Animal products: sheep & goat	Kidney, edible offals	0.01	0.8	
	Muscle	0.06	0.1	
	Fat	0.1	0.2	
Animal products: poultry	Kidney, edible offals	0.5	0.9	
	Muscle, fat, liver, kidney, edible offals	0.05*	0.03*	
Milk	Cattle, sheep, goat, horse	0.05*	0.06	
Bird eggs	Chicken, duck, geese, quail	0.05*	0.15	

1 Including degradation and reaction products which can be determined as 3-(3-thianyl)glutaric acid S-dioxide (BH 517-TGSQ2) including other mixtures of constituent isomers including cycloxydim-M (sum of isomers).
* Limit of determination.
Shading indicates a decrease in MRL.
Source: based on SANTE/10044/2020, Annex II

www.agrinfo.eu

Source: based on Commission Regulation (EU) 2023/173

Disclaimer: *Under no circumstances shall COLEAD be liable for any loss, damage, liability or expense incurred or suffered that is claimed to have resulted from the use of information available on this website or any link to external sites. The use of the website is at the user's sole risk and responsibility. This information platform was created and maintained with the financial support of the European Union. Its contents do not, however, reflect the views of the European Union.*