Maximum residue levels for cycloxydim

Published by AGRINFO on 30 Nov 2022; Revised 17 Apr 2025

Commission Regulation (EU) 2025/581 as regards maximum residue levels for cycloxydim, dichlorprop-P, flupyradifurone, methyl nonyl ketone, plant oils/citronella oil, potassium sorbate and potassium phosphonate in or on certain products

What is changing and why?

The maximum residue levels (MRLs) for cycloxydim have been increased as summarised in Table 1.

Following a request for modification of the MRLs for cycloxydim on some products, the European Food Safety Authority has concluded the data are sufficient to support the proposed MRLs for pome fruits, peas, maize/corn, and sugar beet roots.

Timeline

The new MRLs apply from **17 April 2025**.

For more information see the full record on the AGRINFO website – where you can also view the latest AGRINFO Update newsletters and search the database.

Tables & Figures

Table 1 Changes to maximum residue levels for cycloxydim			
Food category	Products	Cycloxydim (mg/kg)	
		Old MRL	New MRL
Pome fruits	Pears	0.09*	0.4
	Quinces	0.09*	0.4
	Medlars	0.09*	0.4
	Loquats/Japanese medlars	0.09*	0.4
Legume vegetables	Peas (with pods)	2	9
Cereals	Maize/corn	0.2	0.3
Sugar plants	Sugar beet roots	0.2	0.3
* Limit of determination.			

www.agrinfo.eu

Source: based on Commission Regulation 2025/581

Disclaimer: *Under no circumstances shall COLEAD be liable for any loss, damage, liability or expense incurred or suffered that is claimed to have resulted from the use of information available on this website or any link to external sites. The use of the website is at the user's sole risk and responsibility. This information platform was created and maintained with the financial support of the European Union. Its contents do not, however, reflect the views of the European Union.*