

## Maximum residue levels for epoxiconazole



*Published by AGRINFO on 30 Jan 2023*

For more information see the [full record](#) on the AGRINFO website – where you can also view the latest [AGRINFO Update](#) newsletters and [search](#) the database.

## Tables & Figures

Table 1 Maximum residue levels for epoxiconazole			
Food category	Products	Epoxiconazole (mg/kg)	
		Old MRL	New MRL
Miscellaneous fruits	Bananas	0.5	0.01*
Brassica vegetables	Head cabbages	0.2	0.01*
Legume vegetables	Peas (with pods)	0.1	0.01*
Pulses	Peas	0.1	0.01*
Cereals	Barley, oat	1.5	0.01*
	Buckwheat, maize/ corn, common millet/ proso millet, rice, sorghum	0.1	0.01*
	Rye, wheat	0.6	0.01*
Hops	Hops	0.1	0.05*
Spices	All seed spices, fruit spices, bark spices, roots and rhizomes spices, bud spices, flower pistil spices, aril spices	0.1	0.05*
Sugar plants	Sugar beet roots	0.1	0.01*
	Chicory roots	0.2	0.01*
Products of animal origin	Swine/ bovine/ sheep/ goat/ equine/ other farmed terrestrial animals: Liver	0.2	0.01*
	Swine edible offals	0.05	0.01*
	Bovine/ sheep/ goat/ equine/ other farmed terrestrial animals: Kidney, edible offals	0.02	0.01*

\* Limit of determination.  
Source: based on [SANTE/10176/2022 rev.2](#)



  
[www.agrinfo.eu](http://www.agrinfo.eu)

Source: based on Commission Regulation (EU) 2023/128

**Disclaimer:** Under no circumstances shall COLEAD be liable for any loss, damage, liability or expense incurred or suffered that is claimed to have resulted from the use of information

*available on this website or any link to external sites. The use of the website is at the user's sole risk and responsibility. This information platform was created and maintained with the financial support of the European Union. Its contents do not, however, reflect the views of the European Union.*