

Maximum residue levels for etridiazole

Published by AGRINFO on 01 Aug 2023; Revised 05 Feb 2024

Commission Regulation (EU) [2024/345](#) as regards maximum residue levels for desmedipham, etridiazole, flurtamone, profoxydim, difenacoum and potassium permanganate in or on certain products

What is changing and why?

The EU has reduced the maximum residue levels (MRLs) for etridiazole to the limit of determination for all products (see Table 1). In addition, where advances in analytical techniques allow lower limits of determination (LODs) to be achieved, the LODs will be reduced accordingly. (The LOD is the lowest level that can be detected using the most modern and reliable analytical methods.)

Actions

For the products in Table 1, suppliers should explore alternative solutions to their current use of etridiazole.

Timeline

The new MRLs will apply from **12 August 2024**.

For more information see the [full record](#) on the AGRINFO website – where you can also view the latest [AGRINFO Update](#) newsletters and [search](#) the database.

Tables & Figures

Food category	Products	Etridiazole (mg/kg)	
		Old MRL	New MRL
Berries and small fruits	Strawberries	0.1	0.01*
Fruiting vegetables	Sweet peppers/ bell peppers	0.1	0.01*
Cucurbits	Cucumbers, gherkins, courgettes	0.4	0.01*
Oilseeds	Cotton seeds	0.1	0.01*

* Limit of determination.



www.agrinfo.eu

Source: based on Regulation (EU) [2024/345](#)

Disclaimer: Under no circumstances shall COLEAD be liable for any loss, damage, liability or expense incurred or suffered that is claimed to have resulted from the use of information available on this website or any link to external sites. The use of the website is at the user's sole risk and responsibility. This information platform was created and maintained with the financial support of the European Union. Its contents do not, however, reflect the views of the European Union.