


Maximum residue levels for fenpicoxamid

Published by AGRINFO on 29 Jun 2023

For more information see the [full record](#) on the AGRINFO website – where you can also view the latest [AGRINFO Update](#) newsletters and [search](#) the database.

Tables & Figures

Table 1 Changes to maximum residue levels for fenpicoxamid			
Food category	Products	Fenpicoxamid (mg/kg)	
		Old MRL	New MRL
Animal products	Muscle, fat from swine, cattle, sheep, goats, horses, other farmed animals	0.01*	0.015
	Kidney from swine, goats, horses, other farmed animals	0.01*	0.02
	Liver from swine, cattle, goats, horses, other farmed animals	0.01*	0.02
	Edible offals from swine, cattle, sheep, goats, horses, other farmed animals	0.01*	0.02
	Milk from cattle, sheep, goats, horses	0.01*	0.015
* Limit of determination.			
 www.agrinfo.eu			

Source: based on Regulation [2023/1069](#)

Disclaimer: *Under no circumstances shall COLEAD be liable for any loss, damage, liability or expense incurred or suffered that is claimed to have resulted from the use of information available on this website or any link to external sites. The use of the website is at the user's sole risk and responsibility. This information platform was created and maintained with the financial support of the European Union. Its contents do not, however, reflect the views of the European Union.*