


# Maximum residue levels for fenpropimorph

Published by AGRINFO on 18 Jan 2023; Revised 11 Apr 2023

For more information see the [full record](#) on the AGRINFO website – where you can also view the latest [AGRINFO Update](#) newsletters and [search](#) the database.

## Tables & Figures

Table 1 Changes to maximum residue levels for fenpropimorph			
Food category	Products	Fenpropimorph (mg/kg)	
		Old MRL	New MRL
Berries and small fruits	Dewberries, raspberries	1.5	0.01*
	Blueberries, cranberries, currants, gooseberries	0.9	0.01*
Root and tuber vegetables	Carrots, horseradishes, parsnips, parsley roots/ Hamburg, salsifies	0.04	0.01*
Cereals	Barley, oat	0.4	0.2
	Rye, wheat	0.15	0.07
Animal products: Swine	Muscle	0.04	0.03
Animal products: Bovine, sheep, goat and equine	Muscle	0.15	0.03
	Fat	0.2	0.05
	Liver	3	0.7
	Edible offals	3	0.7
Animal products	Milk (cattle, sheep, goat, horse, others)	0.015	0.01
Animal products	Wild terrestrial vertebrates	0.04	0.01*
* Limit of determination.			
 www.agrinfo.eu			

Source: based on Regulation (EU) [2023/710](#)

**Disclaimer:** *Under no circumstances shall COLEAD be liable for any loss, damage, liability or expense incurred or suffered that is claimed to have resulted from the use of information available on this website or any link to external sites. The use of the website is at the user's sole risk and responsibility. This information platform was created and maintained with the financial support of the European Union. Its contents do not, however, reflect the views of the European Union.*