

Maximum residue levels for fluxapyroxad

Published by AGRINFO on 23 Jan 2025

Commission Regulation (EU) [2025/115](#) as regards maximum residue levels for fluxapyroxad, lambda-cyhalothrin, metalaxyl, and nicotine in or on certain products

What is changing and why?


The European Union (EU) has raised the maximum residue levels (MRLs) for fluxapyroxad on kaki/Japanese persimmons from 0.01 to 0.2 mg/kg, and on cultivated fungi from 0.01 to 0.3 mg/kg (see Table 1). This follows an application to the EU for modification of these MRLs. An evaluation by the European Food Safety Authority concluded that there are no risks to consumers at these levels.

Timeline

The new MRLs apply from **11 February 2025**.

For more information see the [full record](#) on the AGRINFO website – where you can also view the latest [AGRINFO Update](#) newsletters and [search](#) the database.

Tables & Figures

Table 1 Changes to maximum residue levels for fluxapyroxad			
Food category	Products	Fluxapyroxad (mg/kg)	
		Old MRL	New MRL
Miscellaneous fruits	Kaki/Japanese persimmons	0.01*	0.2
Fungi, mosses, and lichens	Cultivated fungi	0.01*	0.3
* Limit of determination.			
 www.agrininfo.eu			

Source: Based on Regulation [2025/115](#)

Disclaimer: *Under no circumstances shall COLEAD be liable for any loss, damage, liability or expense incurred or suffered that is claimed to have resulted from the use of information available on this website or any link to external sites. The use of the website is at the user's sole risk and responsibility. This information platform was created and maintained with the financial support of the European Union. Its contents do not, however, reflect the views of the European Union.*