

# Maximum residue levels for fuberidazole

*Published by AGRINFO on 09 Jan 2025; Revised 17 Jul 2025*

Commission Regulation (EU) [2025/1163](#) as regards maximum residue levels for chlorpropham, fuberidazole, ipconazole, methoxyfenozide, S-metolachlor and triflusulfuron in or on certain products

## What is changing and why?

Fuberidazole is no longer permitted for use in the European Union, therefore the EU decided to set all maximum residue levels (MRLs) for fuberidazole at the product-specific limits of determination (LODs). The LOD on some cereals (wheat, rye, oats, and barley) are lowered from 0.05 to 0.01 mg/kg. (The LOD is the lowest level that can be detected using the most modern and reliable analytical methods.)

## Actions

Laboratories testing for fuberidazole in products for export to the EU market should verify that their analytical methods are able to test at the proposed new LODs.

## Timeline

The new MRLs apply from **6 January 2026**.

For more information see the [full record](#) on the AGRINFO website – where you can also view the latest [AGRINFO Update](#) newsletters and [search](#) the database.

**Disclaimer:** *Under no circumstances shall COLEAD be liable for any loss, damage, liability or expense incurred or suffered that is claimed to have resulted from the use of information available on this website or any link to external sites. The use of the website is at the user's sole risk and responsibility. This information platform was created and maintained with the financial support of the European Union. Its contents do not, however, reflect the views of the European Union.*