

Maximum residue levels for methoxyfenozide


Published by AGRINFO on 30 Nov 2022; Revised 28 Jun 2023

For more information see the [full record](#) on the AGRINFO website – where you can also view the latest [AGRINFO Update](#) newsletters and [search](#) the database.

Tables & Figures

Table 1 Changes to maximum residue levels for methoxyfenozide			
Food category	Products	Methoxyfenozide (mg/kg)	
		Old MRL	New MRL
Pome fruits	Apples, pears	2	0.01*
Stone fruits	Peaches	2	0.01*
Brassica vegetables	Broccoli	2	0.01*
Fruiting vegetables	Tomatoes	2	0.6
Leaf vegetables, herbs and edible flowers	Basil, edible flowers	4	80
Teas	Teas	0.05*	80
Coffee beans	Coffee beans	0.05*	0.15
Sugar plants	Sugar canes	0.01*	0.015

* Limit of determination.



www.agrininfo.eu

Source: based on Regulations [2023/1069](#) and [2023/1069](#)

Disclaimer: *Under no circumstances shall COLEAD be liable for any loss, damage, liability or expense incurred or suffered that is claimed to have resulted from the use of information available on this website or any link to external sites. The use of the website is at the user's sole risk and responsibility. This information platform was created and maintained with the financial support of the European Union. Its contents do not, however, reflect the views of the European*

Union.