

Maximum residue levels for methyl nonyl ketone

Published by AGRINFO on 01 Jul 2023; Revised 19 Apr 2024

Regulation

Regulation [2024/352](#) as regards maximum residue levels for (Z)-13-hexadecen-11-yn-1-yl acetate, (Z,Z,Z,Z)-7,13,16,19-docosatetraen-1-yl isobutyrate, acrinathrin, azimsulfuron, famoxadone, prochloraz and sodium hypochlorite in or on certain products

What is changing and why?

The European Commission has set the maximum residue levels (MRLs) for methyl nonyl ketone to the LOD of 0.01 mg/kg on all products. Since 26 May 2017, methyl nonyl ketone is no longer permitted for use in the EU. (The LOD is the lowest level that can be detected using the most modern and reliable analytical methods.)

Timeline

The new MRLs will apply from **12 August 2024**.

For more information see the [full record](#) on the AGRINFO website – where you can also view the latest [AGRINFO Update](#) newsletters and [search](#) the database.

Disclaimer: *Under no circumstances shall COLEAD be liable for any loss, damage, liability or expense incurred or suffered that is claimed to have resulted from the use of information available on this website or any link to external sites. The use of the website is at the user's sole risk and responsibility. This information platform was created and maintained with the financial support of the European Union. Its contents do not, however, reflect the views of the European Union.*