

Maximum residue levels for metolachlor

Published by AGRINFO on 28 Dec 2024; Revised 20 Jun 2025

EU lowers maximum residue levels for metolachlor

Commission Regulation (EU) [2025/1163](#) of 13 June 2025 amending Annexes II, III and V to Regulation (EC) No 396/2005 of the European Parliament and of the Council as regards maximum residue levels for chlorpropham, fuberidazole, ipconazole, methoxyfenozide, S-metolachlor and triflurosulfuron in or on certain products

Update

The European Commission has lowered the maximum residue levels (MRLs) for metolachlor to 0.01 mg/kg for most products.

Impacted products

Algae/prokaryotes, almonds, American persimmons/Virginia kaki, apples, apricots, arrowroots, asparagus, aubergines/eggplants, avocados, azaroles/Mediterranean medlars, baby leaf crops (incl. brassicas), bamboo shoots, bananas, barley, basil and edible flowers, beans (dry, with pods, without pods), beetroots, blackberries, blueberries, borage seeds, Brazil nuts, breadfruits, broccoli, Brussels sprouts, buckwheat/other pseudocereals, carambolas, cardoons, carrots, cashew nuts, cassava roots/manioc, castor beans, cauliflowers, celeriacs/turnip rooted celeries, celeries, celery leaves, chards/beet leaves, cherimoyas, cherries (sweet), chervil, chestnuts, chicory roots, Chinese cabbages/pe-tsai, chives, coconuts, common millet/proso millet, cotton seeds, courgettes, cranberries, cresses/other sprouts and shoots, cucumbers, currants (black, red, white), dates, dewberries, durians, elderberries, escaroles/broad-leaved endives, figs, Florence fennels, fungi, garlic, gherkins, globe artichokes, gold of pleasure seeds, gooseberries (green, red, yellow), granate apples/pomegranates, grape leaves, grapefruits, guavas, hazelnuts/cobnuts, head cabbages, hemp seeds, horseradishes, jambuls/jambolans, Jerusalem artichokes, kaki/Japanese persimmons, kales, kapok, kiwi fruits (green, red, yellow), kohlrabies, kumquats, lamb's lettuces/corn salads, land cresses, laurel/bay leaves, leeks, lemons, lentils, lettuces, limes, linseeds, lychees, loquats/Japanese medlars, lupins/lupini beans, macadamias, maize/corn, mandarins, mangoes, medlars, melons, mosses/lichens, mulberries (black, white), mustard seeds, oats, oil palm fruits, oil palm kernels, okra/lady's fingers, olives for oil production, onions, oranges, palm hearts, papayas, parsley, parsley roots/Hamburg root parsley, parsnips, passionfruits/maracujas, peaches, peanuts/groundnuts, pears, peas (dry, with pods, without pods), pecans, pine nut kernels, pineapples, pistachios, plums, poppy seeds, potatoes, prickly pears/cactus fruits, pumpkin seeds, pumpkins, purslanes, quinces, radishes, rapeseeds/canola seeds, raspberries (red, yellow), red mustards, rhubarbs, rice, Roman rocket/rucola, rose hips, rosemary, rye, safflower seeds, sage, salsifies, sesame seeds, shallots, sorghum, soursops/guanabanas, soyabeans, spinaches, spring onions/green onions/Welsh onions, star apples/cainitos, strawberries, sugar beet roots, sugar canes, sunflower seeds, swedes/rutabagas, sweet corn, sweet peppers/bell peppers, sweet potatoes, table grapes, table olives, tarragon, thyme, tomatoes, turnips, walnuts, watercresses, watermelons, wheat, wild fungi, wine grapes, witloofs/Belgian endives, yams

What is changing?

The MRLs for metolachlor are currently set at the limit of determination (LOD, the lowest level that can be detected using the most modern and reliable analytical methods) between 0.01 and 0.05 mg/kg. (The LOD is the the lowest level that can be detected using the most modern and reliable analytical methods.) The European Commission has reduced the MRL (LOD) for metolachlor to 0.01 mg/kg on all fruits, vegetables, pulses, oilseeds, cereals, and sugar (see "Impacted products"). For herbs, the new MRL will be 0.02 mg/kg.

The residue definition of metolachlor is “Metolachlor and S-metolachlor [metolachlor including other mixtures of constituent isomers including S-metolachlor (sum of isomers)]” (Regulation [2025/1163](#), Annex V).

Why?

The approval for use of S-metolachlor expired in January 2024 (see [S-metolachlor: Non renewal of EU approval](#)). As no Codex MRLs or import tolerances exist for this substance, the MRLs are set at the LOD, and these LODs have been updated to reflect the most recent available analytical methods.

Timeline

The new MRLs apply from **6 January 2026**.

What are the major implications for exporting countries?

As the MRLs were already set at the LOD, the proposed changes are not expected to have a significant impact on the use of this substance on products intended for the EU market. However, these changes may have an impact on the analytical methods used by laboratories.

Recommended Actions

Laboratories testing for metolachlor in products for export to the EU market should verify that their analytical methods are able to test at the proposed new LODs.

Background

MRLs are set in accordance with the rules set out in Regulation [396/2005](#). For information on current MRLs for other substances, please consult the [EU Pesticide Residues database](#).

Sources

Commission Regulation (EU) [2025/1163](#) as regards maximum residue levels for chlorpropham, fuberidazole, ipconazole, methoxyfenozide, S-metolachlor and triflurosulfuron in or on certain products

Visit the [AGRINFO website](#) to view the latest AGRINFO Update newsletters and [search](#) the database.

Disclaimer: *Under no circumstances shall COLEAD be liable for any loss, damage, liability or expense incurred or suffered that is claimed to have resulted from the use of information available on this website or any link to external sites. The use of the website is at the user's sole risk and responsibility. This information platform was created and maintained with the financial support of the European Union. Its contents do not, however, reflect the views of the European Union.*