Maximum residue levels for pinoxaden

Published by AGRINFO on 24 Jun 2026; Revised 31 Jul 2026

Draft Commission Regulation as regards maximum residue levels for acequinocyl, deltamethrin, dodine, maleic hydrazide, pinoxaden and prothioconazole in or on certain products

Draft Annex [PLAN/2025/3053]

What is changing and why?

The European Union (EU) proposes to increase the maximum residue level (MRL) for pinoxaden on barley, rye, and wheat from 0.7 to 0.8 mg/kg. An application for a change of MRL by the producer of pinoxaden was evaluated by the European Food Safety Authority, and the requested changes were found to be safe for consumers.

Timeline

The Regulation is expected to be published in October 2026, and will apply 6 months after publication.

For more information see the full record on the AGRINFO website – where you can also view the latest AGRINFO Update newsletters and search the database.

Disclaimer: *Under no circumstances shall COLEAD be liable for any loss, damage, liability or expense incurred or suffered that is claimed to have resulted from the use of information available on this website or any link to external sites. The use of the website is at the user's sole risk and responsibility. This information platform was created and maintained with the financial support of the European Union. Its contents do not, however, reflect the views of the European Union.*

Copyright © **COLEAD 2026,** *AGRINFO is funded by the European Union and implemented by COLEAD.*