

Maximum residue levels for pyraclostrobin

Published by AGRINFO on 29 Jan 2024; Revised 20 Feb 2026

Commission Regulation (EU) [2026/140](#) as regards maximum residue levels for acequinocyl, chlormequat, metalaxyl-M, pyraclostrobin, sulfoxaflor and trifloxystrobin in or on certain products

Commission Regulation (EU) [2024/342](#) as regards maximum residue levels for cyflumetofen, oxathiapiprolin and pyraclostrobin in or on certain products

What is changing and why?

The European Union (EU) has raised the maximum residue level (MRL) for pyraclostrobin on sweet corn from 0.04 to 0.09 mg/kg, a level that the European Food Safety Authority considers to be safe.

Timeline

The new MRL for sweet corn applies from **11 February 2026**.

For more information see the [full record](#) on the AGRINFO website – where you can also view the latest [AGRINFO Update](#) newsletters and [search](#) the database.

Disclaimer: *Under no circumstances shall COLEAD be liable for any loss, damage, liability or expense incurred or suffered that is claimed to have resulted from the use of information available on this website or any link to external sites. The use of the website is at the user's sole risk and responsibility. This information platform was created and maintained with the financial support of the European Union. Its contents do not, however, reflect the views of the European Union.*