

Maximum residue levels for quinmerac


Published by AGRINFO on 30 Nov 2022

For more information see the [full record](#) on the AGRINFO website – where you can also view the latest [AGRINFO Update](#) newsletters and [search](#) the database.

Tables & Figures

Table 1 Maximum residue levels for quinmerac			
Food category	Products	Quinmerac ¹ (mg/kg)	
		Old MRL	New MRL
Leaf vegetables, herbs and edible flowers	Spinaches, chards/ beet leaves, chervil, chives, celery leaves, parsley, sage, rosemary, thyme, basil and edible flowers, laurel/ bay leaves, tarragon	0.5	0.1*
Sugar plants	Sugar beet roots	0.5	0.15*
Products of animal origin	Bovine: liver, edible offals	0.05*	0.07
	Bovine: kidney	0.05*	0.05
	Sheep: liver/kidney/edible offals	0.05*	0.05
	Goat: liver/kidney/edible offals	0.05*	0.05
	Equine: liver, edible offals	0.05*	0.07
	Equine: kidney	0.05*	0.05

1 Sum of quinmerac and its metabolites BH 518-2 and BH 518-4 expressed as quinmerac.
 * Limit of determination.
 Source: based on Commission Regulation (EU) 2022/1321



 www.agrinfo.eu

Disclaimer: Under no circumstances shall COLEAD be liable for any loss, damage, liability or expense incurred or suffered that is claimed to have resulted from the use of information

available on this website or any link to external sites. The use of the website is at the user's sole risk and responsibility. This information platform was created and maintained with the financial support of the European Union. Its contents do not, however, reflect the views of the European Union.