Maximum residue levels for sulfoxaflor

Published by AGRINFO on 04 Feb 2026

Commission Regulation (EU) 2026/140 as regards maximum residue levels for acequinocyl, chlormequat, metalaxyl-M, pyraclostrobin, sulfoxaflor and trifloxystrobin in or on certain products

What is changing and why?

The European Union (EU) has increased the maximum residue levels (MRLs) for sulfoxaflor as summarised in Table 1.

Timeline

The new MRLs apply from **11 February 2026**.

For more information see the full record on the AGRINFO website – where you can also view the latest AGRINFO Update newsletters and search the database.

Copyright © **COLEAD 2026,** *AGRINFO is funded by the European Union and implemented by COLEAD.*

Tables & Figures

Table 1
Changes to maximum residue levels for sulfoxaflor

Food category	Products	Sulfoxaflor (mg/kg)	
		Previous MRL	New MRL
Fruiting vegetables	Okra/lady's fingers	0.01*	0.07
Leaf vegetables	Lamb's lettuces/corn salads, escaroles/ broad-leaved endives, cresses and other sprouts and shoots, land cresses, Roman rocket/rucola, red mustards, baby leaf crops, purslanes, chards/beet leaves, watercresses	0.01*	0.7
Herbs and edible flowers	Chervil, chives, parsley, sage, rosemary, thyme, basil and edible flowers, laurel/bay leaves, tarragon	0.02*	0.7

* Limit of determination.

AGRINFO
www.agrinfo.eu

Source: based on Regulation (EU) 2026/140

Disclaimer: *Under no circumstances shall COLEAD be liable for any loss, damage, liability or expense incurred or suffered that is claimed to have resulted from the use of information available on this website or any link to external sites. The use of the website is at the user's sole risk and responsibility. This information platform was created and maintained with the financial support of the European Union. Its contents do not, however, reflect the views of the European Union.*

Copyright © **COLEAD 2026,** *AGRINFO is funded by the European Union and implemented by COLEAD.*