

# Maximum residue levels for trifloxystrobin

*Published by AGRINFO on 03 Dec 2023; Revised 02 Feb 2026*

Commission Regulation (EU) [2026/140](#) as regards maximum residue levels for acequinocyl, chlormequat, metalaxyl-M, pyraclostrobin, sulfoxaflor and trifloxystrobin in or on certain products

Commission Regulation (EU) [2024/1342](#) as regards maximum residue levels for deltamethrin, metalaxyl, thiabendazole and trifloxystrobin in or on certain products

## What is changing and why?

The European Union (EU) has raised the maximum residue levels (MRLs) for trifloxystrobin on several products, as summarised in Table 1. This follows an application to modify the MRLs on olives, celeries, leeks, globe artichokes, herbal infusions and seed spices, as well as a request for import tolerance on linseeds. Both applications were assessed and considered safe by the European Food Safety Authority.

In 2024, the EU amended the MRLs for trifloxystrobin as summarised in Table 2.

## Actions

Suppliers of affected products should review their current use of trifloxystrobin and assess whether any changes will be needed to existing good agricultural practices (GAP), or start looking for possible alternative solutions.

## Timeline

The new MRLs in Table 1 apply from **11 February 2026**.

The MRLs in Table 2 applied from 11 December 2024.

For more information see the [full record](#) on the AGRINFO website – where you can also view the latest [AGRINFO Update](#) newsletters and [search](#) the database.

## Tables & Figures

Table 1 Changes to maximum residue levels for trifloxystrobin			
Food category	Products	Trifloxystrobin (mg/kg)	
		Previous MRL	New MRL
Miscellaneous fruit	Table olives	0.3	0.5
Stem vegetables	Celeries	1	15
	Globe artichokes	0.3	0.5
	Leeks	0.7	0.9
Oilseeds	Linseeds	0.01*	0.4
Oil fruits	Olives for oil production	0.3	0.5
Herbal infusions	Chamomile, hibiscus/roselle, rose, jasmine, lime/linden, other flower infusions, strawberry, rooibos, mate/maté, other leaves/ herb infusions	0.05*	30
Spices	Anise/aniseed, black caraway/black cumin, celery, coriander, cumin, dill, fennel, fenugreek, nutmeg, other seed spices	0.05*	50
* Limit of determination.			
 www.agrinfo.eu			

Source: based on Regulation (EU) [2026/140](#)

Table 2 Changes to maximum residue levels for trifloxystrobin			
Food category	Products	Trifloxystrobin (mg/kg)	
		Previous MRL	New MRL
Miscellaneous fruits	Passionfruits/maracujas	4	0.01*
Fruiting vegetables	Sweet peppers/bell peppers	0.4	0.9
Brassica vegetables	Chinese cabbages/pe-tsai	3	0.01*
	Kales	3	2
Leaf vegetables	Chervil, chives, celery leaves, parsley, sage, rosemary, thyme, basil and edible flowers, laurel/bay leaves, tarragon	15	30
Legume vegetables	Beans (with pods)	1	0.8
Cereals	Oats	0.4	0.3
Sugar plants	Chicory roots	0.01*	0.02
Products of animal origin	Honey	0.05*	0.07
* Limit of determination.			
 www.agrinfo.eu			

Source: based on Regulation (EU) [2024/1342](#)

**Disclaimer:** *Under no circumstances shall COLEAD be liable for any loss, damage, liability or expense incurred or suffered that is claimed to have resulted from the use of information available on this website or any link to external sites. The use of the website is at the user's sole risk and responsibility. This information platform was created and maintained with the financial support of the European Union. Its contents do not, however, reflect the views of the European Union.*