


Mercury maximum levels

Published by AGRINFO on 22 Nov 2022

For more information see the [full record](#) on the AGRINFO website – where you can also view the latest [AGRINFO Update](#) newsletters and [search](#) the database.

Tables & Figures

Table 1 Mercury maximum levels		
Product	Maximum levels (mg/kg wet weight)	
	New	Previous
Fishery products and muscle meat of fish, excluding species listed in the rows below.*	0.50	0.50
Muscle meat of the following fish: Axillary seabream (<i>Pagellus acarne</i>) Black scabbardfish (<i>Aphanopus carbo</i>) Blackspot seabream (<i>Pagellus bogaraveo</i>) Bonito (<i>Sarda sarda</i>) Common pandora (<i>Pagellus erythrinus</i>) Escolar (<i>Lepidocybium flavobrunneum</i>) Halibut (<i>Hippoglossus</i> species) Kingklip (<i>Genypterus capensis</i>) Marlin (<i>Makaira</i> species) Megrim (<i>Lepidorhombus</i> species) Oilfish (<i>Ruvettus pretiosus</i>) Orange roughy (<i>Hoplostethus atlanticus</i>) Pink cusk-eel (<i>Genypterus blacodes</i>) Pike (<i>Esox</i> species) Plain bonito (<i>Orcynopsis unicolor</i>) Poor cod (<i>Tricopterus</i> species) Red mullet (<i>Mullus barbatus barbatus</i>) Roundnose grenadier (<i>Coryphaenoides rupestris</i>) Sail fish (<i>Istiophorus</i> species) Silver scabbardfish (<i>Lepidopus caudatus</i>) Snake mackerel (<i>Gempylus serpens</i>) Sturgeon (<i>Acipenser</i> species) Surmullet (<i>Mullus surmuletus</i>) Tuna (<i>Thunnus</i> species, <i>Euthynnus</i> species, <i>Katsuwonus pelamis</i>) Shark (all species) Swordfish (<i>Xiphias gladius</i>)	1.0	**
Cephalopods Marine gastropods Muscle meat of the following fish: Anchovy (<i>Engraulis</i> species) Alaska pollock (<i>Theragra chalcogrammus</i>) Atlantic cod (<i>Gadus morhua</i>) Atlantic herring (<i>Clupea harengus</i>) Basa (<i>Pangasius bocourti</i>) Carp (species belonging to the Cyprinidae family) Common dab (<i>Limanda limanda</i>) Mackerel (<i>Scomber</i> species) European flounder (<i>Platichthys flesus</i>) European plaice (<i>Pleuronectes platessa</i>) European sprat (<i>Sprattus sprattus</i>) Mekong giant catfish (<i>Pangasianodon gigas</i>) Pollock (<i>Pollachius pollachius</i>) Saithe (<i>Pollachius virens</i>) Salmon and trout (<i>Salmo</i> species and <i>Oncorhynchus</i> species, except <i>Salmo trutta</i>) Sardine or pilchard (<i>Dussumieria</i> species, <i>Sardina</i> species, <i>Sardinella</i> species and <i>Sardinops</i> species) Sole (<i>Solea solea</i>) Striped catfish (<i>Pangasianodon hypophthalmus</i>) Whiting (<i>Merlangius merlangus</i>)	0.30	0.50
Salt	0.10	No limit
* The maximum level for crustaceans applies to muscle meat from appendages and abdomen. In the case of crabs and crab-like crustaceans (Brachyura and Anomura), it applies to muscle meat from appendages. ** For fish in bold, no change of maximum level. The other fish listed were previously subject to the general maximum level of 0.5 mg/kg. Source: Commission Regulation (EU) 2022/617, Annex		
 www.agrinfo.eu		

Disclaimer: *Under no circumstances shall COLEAD be liable for any loss, damage, liability or expense incurred or suffered that is claimed to have resulted from the use of information available on this website or any link to external sites. The use of the website is at the user's sole risk and responsibility. This information platform was created and maintained with the financial support of the European Union. Its contents do not, however, reflect the views of the European Union.*