

MRLs for sodium salicylate in poultry

Published by AGRINFO on 20 Nov 2023; Revised 02 Apr 2024

Commission Implementing Regulation (EU) [2024/859](#) as regards the classification of the substance sodium salicylate with respect to its maximum residue limit in foodstuffs of animal origin

What is changing and why?

The EU has now set the following maximum residue levels (MRLs) for sodium salicylate in tissues of poultry (other than turkey):

- muscle, and skin and fat in natural proportion: 250 µg/kg
- liver: 500 µg/kg
- kidney: 1000 µg/kg.

MRLs for sodium salicylate in turkey were already set under Regulation [37/2010](#).

Sodium salicylate may not be used in poultry that is producing eggs for human consumption, due to a lack of data on egg consumption.

Timeline

The new MRLs will apply from **8 April 2024**.

For more information see the [full record](#) on the AGRINFO website – where you can also view the latest [AGRINFO Update](#) newsletters and [search](#) the database.

Disclaimer: *Under no circumstances shall COLEAD be liable for any loss, damage, liability or expense incurred or suffered that is claimed to have resulted from the use of information available on this website or any link to external sites. The use of the website is at the user's sole risk and responsibility. This information platform was created and maintained with the financial support of the European Union. Its contents do not, however, reflect the views of the European Union.*