

Maximum residue levels for chlorpropham

Published by AGRINFO on 09 Jan 2025; Revised 18 Jun 2025

Commission Regulation (EU) [2025/1163](#) as regards maximum residue levels for chlorpropham, fuberidazole, ipconazole, methoxyfenozide, S-metolachlor and triflusaluron in or on certain products

What is changing and why?

Chlorpropham is not approved in the European Union (EU), and most maximum residue levels (MRLs) are set at the product-specific limits of determination (LOD, the lowest level that can be detected using the most modern and reliable analytical methods).

One exception is potatoes. The EU has now reduced the MRL for chlorpropham in potatoes from 0.35 to 0.2 mg/kg.

Actions

Suppliers of potatoes to the EU market should verify current good agricultural practices (GAP) to ensure compliance with a reduced MRL. It is also important to continue monitoring chlorpropham levels in potatoes so that data can be submitted to the European Commission before approximately July 2027, when a new review will be carried out.

Timeline

The new MRL applies from **6 January 2026**.

For more information see the [full record](#) on the AGRINFO website – where you can also view the latest [AGRINFO Update](#) newsletters and [search](#) the database.

Disclaimer: *Under no circumstances shall COLEAD be liable for any loss, damage, liability or expense incurred or suffered that is claimed to have resulted from the use of information available on this website or any link to external sites. The use of the website is at the user's sole risk and responsibility. This information platform was created and maintained with the financial support of the European Union. Its contents do not, however, reflect the views of the European Union.*